

Crossing probabilities for critical Bernoulli percolation on slabs

Deepan Basu *

Artem Sapozhnikov †

March 23, 2016

Abstract

We prove that in the critical Bernoulli percolation on two dimensional lattice slabs the probabilities of open left-right crossings of rectangles with any given aspect ratio are uniformly positive.

1 Introduction

One of the main tools in the study of planar percolation models at criticality is the Russo-Seymour-Welsh (RSW) theorem. It states that the probability that an open path connects the left and right sides of a rectangle is bounded away from 0 and 1 by constants that only depend on the aspect ratio of the rectangle. This theorem was first proved for critical Bernoulli percolation on planar lattices in [9, 11, 10, 7] and recently has been extended to some other planar models, perhaps most notably to the FK-percolation [3, 5] and Voronoi percolation [2, 12].

In this note we consider the critical Bernoulli percolation on two dimensional slabs $\mathbb{Z}^2 \times \{0, \dots, k-1\}^{d-2}$. We prove that the probability of crossing a rectangle is bounded from below by a positive constant which only depends on the aspect ratio of the rectangle and the slab parameter k , but does not depend on the size of the rectangle. Our work is inspired by a recent paper of Duminil-Copin, Sidoravicius, and Tassion [4] where they prove the absence of percolation at criticality for slabs and develop techniques for “glueing” open paths. Our proof is partly based on these new ideas.

2 Notation and result

Fix an integer $k \geq 1$, and define the slab of width k by

$$\mathbb{S} = \mathbb{Z}^2 \times \{0, \dots, k-1\}^{d-2}.$$

We consider Bernoulli bond percolation on $\mathbb{S}$ with parameter $p \in [0, 1]$, and denote the corresponding measure by $\mathbb{P}_p$. Let p_c be the critical threshold for percolation, i.e.,

$$p_c = \inf \{p : \mathbb{P}_p[\text{open connected component of } 0 \text{ in } \mathbb{S} \text{ is infinite}] > 0\},$$

*Max-Planck Institute for Mathematics in the Sciences, Inselstrasse 22, 04103 Leipzig, Germany. email: deepan.basu@mis.mpg.de

†University of Leipzig, Department of Mathematics, Augustusplatz 10, 04109 Leipzig, Germany. email: artem.sapozhnikov@math.uni-leipzig.de

Bibliographic note: After this project was completed, we learned that an alternative proof of our main result was independently obtained by Newman, Tassion, and Wu, which was subsequently written up in [8]. In the same paper, they prove that in the critical Bernoulli percolation on slabs, the probabilities of open left-right crossings of rectangles with any given aspect ratio are uniformly smaller than 1, see Remark 2.2.

and define the measure $\mathbb{P} = \mathbb{P}_{p_c}$.

For a subset A of vertices of $\mathbb{Z}^2$, let

$$\overline{A} = A \times \{0, \dots, k-1\}^{d-2}.$$

Define a rectangle and its left and right boundary regions by

$$B(m, n) = \overline{[0, m] \times [0, n]}, \quad L(m, n) = \overline{\{0\} \times [0, n]}, \quad R(m, n) = \overline{\{m-1\} \times [0, n]}.$$

Consider the crossing event

$$\text{LR}(m, n) = \{L(m, n) \text{ is connected to } R(m, n) \text{ by an open path in } B(m, n)\}$$

and the crossing probability

$$p(m, n) = \mathbb{P}[\text{LR}(m, n)].$$

In this note we prove the following theorem.

Theorem 2.1. *For any $\rho \in (0, \infty)$,*

$$\liminf_{n \rightarrow \infty} p(\lfloor \rho n \rfloor, n) > 0. \quad (2.1)$$

Proof. As some of the “glueing” ideas used in the proof are unnecessary for $k = 1$ and easier for $k \geq 2$, we *assume from now on without further mentioning that $k \geq 2$* . The theorem is proved in 3 steps:

- The result holds for all $\rho \in (0, 1)$. This is well known. We give a proof in Proposition 3.1.
- If the result holds for some $\rho > 1$, then it holds for all $\rho > 1$. This is a well known fact in planar percolation. We prove the slab version in Proposition 5.1 using the planar approach together with a novel technique for glueing paths from [4] (see Lemma 4.2).
- There exist $c > 0$ and $C < \infty$ such that for all $n \geq 1$, $p(44n, 43n) \geq c \cdot p(43n, 44n)^C$. This inequality is the main contribution of this paper. We prove it in Proposition 6.1.

□

Remark 2.2. For $\rho < 1$, the result of Theorem 2.1 holds in any dimension $d \geq 2$. We believe that it also holds for $\rho \geq 1$, but do not know a proof. Our method unfortunately crucially relies on planarity of slabs. If dimension is sufficiently high, it is proved in [1] that the crossing probabilities tend to 1 as $n \rightarrow \infty$. We believe that for percolation on slabs (and in low dimensions) for every $\rho > 0$, $\limsup_{n \rightarrow \infty} p(\lfloor \rho n \rfloor, n) < 1$, but do not have a proof yet.¹

Earlier we defined $\overline{A}$ as a subset of $\mathbb{S}$ for each $A \subset \mathbb{Z}^2$. In the proofs we will often use the same notation $\overline{A}$ for $A \subset \mathbb{S}$ meaning

$$\overline{A} = \{z = (z_1, \dots, z_d) \in \mathbb{S} : (z_1, z_2, x_3, \dots, x_d) \in A \text{ for some } x_3, \dots, x_d\}.$$

This way, for each $A \subset \mathbb{Z}^2$, $\overline{A}$ defined earlier is the same as $\overline{A \times \{0\}^{d-2}}$ defined just above.

¹For slabs, this was recently proved in [8].

3 Crossings of narrow rectangles

The following proposition is an adaptation to slabs of a well known fact about the probabilities of crossing hypercubes of fixed aspect ratio in the easy direction.

Proposition 3.1. *For any $\rho \in (0, 1)$, (2.1) holds.*

Proof. It suffices to prove that for all $L \in \mathbb{N}$, $\liminf_{n \rightarrow \infty} p(n, nL) > 0$. The classical recursive inequality (applied to slabs) states that for every $p \in [0, 1]$, $L, n \geq 1$,

$$\mathbb{P}_p[\text{LR}(2n, 2nL)] \leq ((3L - 1) \cdot \mathbb{P}_p[\text{LR}(n, nL)])^2,$$

which implies that for all $s \geq 0$,

$$\mathbb{P}_p[\text{LR}(2^s n, 2^s nL)] \leq ((3L - 1)^2 \cdot \mathbb{P}_p[\text{LR}(n, nL)])^{2^s}.$$

Fix $L \in \mathbb{N}$. If $\liminf_{n \rightarrow \infty} p(n, nL) = 0$, there exists $n \in \mathbb{N}$ such that $p(n, nL) < \frac{1}{2(3L-1)^2}$. Since the crossing probability $\mathbb{P}_p[\text{LR}(n, nL)]$ is continuous in p , there also exists $p > p_c$ such that $\mathbb{P}_p[\text{LR}(n, nL)] < \frac{1}{2(3L-1)^2}$. For this choice of parameters, $\lim_{s \rightarrow \infty} \mathbb{P}_p[\text{LR}(2^s n, 2^s nL)] = 0$, which is impossible, since for every $p > p_c$, this limit equals to 1 (see, e.g., [6]). Thus, $\liminf_{n \rightarrow \infty} p(n, nL) > 0$. $\square$

4 Glueing

In this section we recall a new technique for glueing paths from [4]. It will be used to adapt some arguments from planar percolation to slabs. We begin with a classical combinatorial lemma about local modifications, see, e.g., [4, Lemma 7].

Lemma 4.1. *Let $n \geq 1$ and $p \in (0, 1)$. Let $A, B \subseteq \{0, 1\}^n$ and $\mathbf{P}_p$ a product measure on $\{0, 1\}^n$ with parameter p , i.e.,*

$$\mathbf{P}_p[\omega] = \prod_{i=1}^n p^{\omega_i} (1-p)^{1-\omega_i}, \quad \omega \in \{0, 1\}^n.$$

If there exists a map $f : A \rightarrow B$ such that for every $\omega' \in B$, there exists a set $S \subseteq \{1, \dots, n\}$ such that $|S| \leq s$ and

$$\omega_i = \omega'_i, \quad \text{for all } i \notin S \text{ and } \omega \in f^{-1}(\omega'),$$

then

$$\mathbf{P}_p[A] \leq \left(\frac{2}{\min(p, 1-p)} \right)^s \cdot \mathbf{P}_p[B].$$

We will often apply Lemma 4.1 in the case $p = p_c$ and s being not bigger than the number of edges in $[-3, 3]^2$. Therefore, we define

$$C_* = \left(\frac{2}{\min(p, 1-p)} \right)^{d \cdot 7^2 \cdot k^{d-2}}, \quad c_* = \frac{1}{1 + C_*}.$$

The following lemma is essentially proven in [4, Lemma 6].

Lemma 4.2. *Let X_1, X_2, Y_1 , and Y_2 be disjoint connected subsets of the interior vertex boundary of $[0, m) \times [0, n)$ arranged in a counter-clockwise order. Then*

$$\mathbb{P}[\overline{X_1} \text{ is connected to } \overline{X_2} \text{ in } B(m, n)] \geq c_* \cdot \mathbb{P} \left[\begin{array}{l} \overline{X_1} \text{ is connected to } \overline{Y_1} \text{ in } B(m, n), \\ \overline{X_2} \text{ is connected to } \overline{Y_2} \text{ in } B(m, n) \end{array} \right]$$

Proof. Let

$$\begin{aligned} E_i &= \{\overline{X_i} \text{ is connected to } \overline{Y_i} \text{ in } B(m, n)\}, \\ X &= \{\overline{X_1} \text{ is connected to } \overline{X_2} \text{ in } B(m, n)\}, \end{aligned}$$

It suffices to prove that $\mathbb{P}[E_1 \cap E_2 \cap X^c] \leq C_* \cdot \mathbb{P}[X]$. We will use Lemma 4.1, where a suitable function $f : E_1 \cap E_2 \cap X^c \rightarrow X$ will be constructed using ideas from [4, Lemma 6].

We fix an order on edges $\{e : |e| = 1\}$ in $\mathbb{Z}^d$ and enumerate all the vertices of $\mathbb{S}$ arbitrarily. Define an order $<$ on self-avoiding paths from $\overline{X_1}$ to $\overline{Y_1}$ in $B(m, n)$ as follows. If $\gamma = (\gamma_0, \dots, \gamma_n)$ and $\gamma' = (\gamma'_0, \dots, \gamma'_{n'})$ are two such paths, then $\gamma < \gamma'$ if

- γ_0 has a smaller number than γ'_0 , or
- $n < n'$ and $\gamma = (\gamma'_0, \dots, \gamma'_n)$, or
- there exists $k < \min(n, n')$ such that $(\gamma_0, \dots, \gamma_k) = (\gamma'_0, \dots, \gamma'_k)$, and the edge $\{0, \gamma_{k+1} - \gamma_k\}$ is smaller than $\{0, \gamma'_{k+1} - \gamma'_k\}$.

For $\omega \in E_1$, let $\gamma_{\min}(\omega)$ be the minimal open self-avoiding path from $\overline{X_1}$ to $\overline{Y_1}$ for the above defined order. Exactly the same construction as in the proof of [4, Lemma 6, Fact 2] gives a function $f : E_1 \cap E_2 \cap X^c \rightarrow E_1 \cap X$ such that for every $\omega \in E_1 \cap E_2 \cap X^c$, there exists a $z \in \gamma_{\min}(\omega)$ such that

- $z \in \gamma_{\min}(f(\omega))$,
- z is a unique vertex on $\gamma_{\min}(f(\omega))$ connected to $\overline{X_2}$ by an open path that does not use edges of $\gamma_{\min}(f(\omega))$,
- $\omega_e = f(\omega)_e$ for all $e \notin \overline{z + [-3, 3]^2 \times \{0\}^{d-2}}$.

Roughly speaking, one chooses $z \in \gamma_{\min}(\omega)$ such that $\overline{z}$ is connected to $\overline{X_2}$ and modifies the configuration in $\overline{z + [-3, 3]^2 \times \{0\}^{d-2}}$ so that in the new configuration z is connected to $\overline{X_1}$, $\overline{Y_1}$, and $\overline{X_2}$ by open paths having only vertex z in common and so that the minimal open path from $\overline{X_1}$ to $\overline{Y_1}$ still passes through z . This tricky construction is very carefully explained in the proof of [4, Lemma 6, Fact 2], therefore we do not repeat it here.

The function f satisfies the conditions of Lemma 4.1 with s being the number of edges in $\overline{[-3, 3]^2}$. Thus, $\mathbb{P}[E_1 \cap E_2 \cap X^c] \leq C_* \cdot \mathbb{P}[E_1 \cap X] \leq C_* \cdot \mathbb{P}[X]$. The proof of the lemma is complete. $\square$

Corollary 4.3. *By Lemma 4.2 and the FKG inequality,*

$$\begin{aligned} \mathbb{P}[\overline{X_1} \text{ is connected to } \overline{X_2} \text{ in } B(m, n)] \\ \geq c_* \cdot \mathbb{P}[\overline{X_1} \text{ is connected to } \overline{Y_1} \text{ in } B(m, n)] \cdot \mathbb{P}[\overline{X_2} \text{ is connected to } \overline{Y_2} \text{ in } B(m, n)]. \end{aligned}$$

5 Crossings of wide rectangles

Proposition 5.1. *If (2.1) holds for some $\rho > 1$, then it holds for all $\rho > 1$.*

Proof. This is immediate from the following inequality, which relates the crossing probability of a long rectangle with that of a shorter one. For all $m > n$,

$$p(2m - n, n) \geq \frac{1}{4} \cdot c_*^3 \cdot p(m, n)^4. \quad (5.1)$$

The inequality (5.1) follows from two applications of Corollary 4.3 illustrated on Figure 1.




Figure 1: (a) left-right crossing of $B(m, n)$ and top-bottom crossing of a $\overline{[m - n, m] \times [0, n]}$ landing on the right half of the bottom, (b) path from $L(m, n)$ to $\overline{[m - \frac{n}{2}, m] \times \{0\}}$ in $B(m, n)$, (c) paths from $L(2m - n, n)$ to $\overline{[m - \frac{n}{2}, m] \times \{0\}}$ and from $\overline{[m - n, m - \frac{n}{2}] \times \{0\}}$ to $R(2m - n, n)$ in $B(2m - n, n)$, (d) left-right crossing of $B(2m - n, n)$.

□

6 Crossings of rectangles: short and long directions

The main contribution of this paper is the following proposition, which relates the crossing probability of a rectangle in the long direction with the one in the short.

Proposition 6.1. *For all $n \in \mathbb{N}$,*

$$p(44n, 43n) \geq \frac{c_*^{21} \cdot p(43n, 44n)^{198}}{10^{154}}. \quad (6.1)$$

Proof. Fix $n \in \mathbb{N}$. We write

$$B = B(43n, 44n), \quad L = L(43n, 44n), \quad R = R(43n, 44n),$$

and define

$$c = p(43n, 44n), \quad c' = \frac{c_*^{21} \cdot c^{198}}{10^{154}}.$$

We prove the proposition by considering several cases. The first 2 steps are inspired by the ideas of Bollobás and Riordan from [2], and aimed at restricting possible shapes of left-right crossings. Steps 3 and 4 contain preliminary estimates needed to implement the main idea in Step 5.

Step 1. We first consider the case when there is a considerable probability that a left-right crossing of B stays away from the top or bottom boundary of B , see Figure 2. Assume that

$$p(43n, 42n) \geq \frac{c}{100}.$$




Figure 2: Left-right crossing staying at least $2n$ away from the top of $B(43n, 44n)$.

Then by (5.1),

$$p(44n, 43n) \geq p(44n, 42n) \geq \frac{1}{4} \cdot c_*^3 \cdot p(43n, 42n)^4 \geq c',$$

which implies (6.1). Thus, we may assume that

$$p(43n, 42n) < \frac{c}{100}. \quad (6.2)$$

Step 2. Next, we consider the case when there is a considerable probability that a left-right crossing of B starts sufficiently far away from the middle of L . Let

$$S = \overline{\{0\} \times [20n, 24n]} \quad (6.3)$$

be the middle of L . Assume that

$$\mathbb{P}[L \setminus S \text{ is connected to } R \text{ in } B] \geq \frac{c}{10}.$$

Then, by reflectional symmetry,

$$\mathbb{P}[\overline{\{0\} \times [24n, 44n]} \text{ is connected to } R \text{ in } B] \geq \frac{c}{20}.$$

By assumption (6.2),

$$\mathbb{P}[\overline{\{0\} \times [24n, 44n]} \text{ is connected to } \overline{[0, 43n) \times \{2n\}} \text{ in } B] \geq \frac{c}{20} - \frac{c}{100} \geq \frac{c}{100}.$$

By rotational symmetry, the above display states precisely that

$$\mathbb{P}[\overline{\{0\} \times [0, 43n)} \text{ is connected to } \overline{[22n, 42n) \times \{0\}} \text{ in } B(42n, 43n)] \geq \frac{c}{100}.$$

Similarly to the second application of Corollary 4.3 in the proof of (5.1), see Figure 3, one gets

$$p(44n, 43n) \geq c_* \cdot \left(\frac{c}{100}\right)^2 \geq c',$$

which is precisely (6.1).




Figure 3: (a) part of L above S is connected to $\overline{[0, 43n] \times \{2n\}}$ in B , (b) rotation of (a) by $\frac{\pi}{2}$, (c) $L(44n, 43n)$ is connected to $\overline{[22n, 42n] \times \{0\}}$ and $\overline{[2n, 22n] \times \{0\}}$ is connected to $R(44n, 43n)$, (d) left-right crossing of $B(44n, 43n)$.

Thus, we may assume, in addition to (6.2), that

$$\mathbb{P}[L \setminus S \text{ is connected to } R \text{ in } B] < \frac{c}{10}. \quad (6.4)$$

Step 3. Here we consider the case when there is a considerable probability that two well-separated subsegments of L are connected. For integers $a < b$, let

$$T_{ab} = \overline{[0, 43n] \times [a, b]} \quad \text{and} \quad T = \overline{[0, 43n] \times \mathbb{Z}}.$$

Assume that for some $a < b$,

$$\mathbb{P}\left[\overline{\{0\} \times [0, 4n]} \text{ is connected to } \overline{\{0\} \times [8n, 12n]} \text{ in } T_{ab}\right] \geq \frac{c_* \cdot c^{18}}{10^{14}}.$$

Then, by repetitive use of Corollary 4.3, see Figure 4, for each $m \geq 1$,

$$\mathbb{P}\left[\overline{\{0\} \times [0, 4n]} \text{ is connected to } \overline{\{0\} \times [4n(m+1), 4n(m+2)]} \text{ in } T\right] \geq \frac{c_*^{2m-1} \cdot c^{18m}}{10^{14m}}.$$




Figure 4: Vertical extension of open paths.

Note that if $m = 11$, then the event on the left hand side implies that there is a vertical crossing of $\overline{[0, 43n)} \times [4n, 48n)$. Thus,

$$p(44n, 43n) \geq \frac{c_*^{21} \cdot c^{198}}{10^{154}} = c',$$

which gives (6.1). Therefore, we may assume, in addition to (6.2) and (6.4), that

$$\mathbb{P} \left[\overline{\{0\} \times [0, 4n)} \text{ is connected to } \overline{\{0\} \times [8n, 12n)} \text{ in } T_{ab} \right] < \frac{c_* \cdot c^{18}}{10^{14}}, \quad \text{for all } a < b. \quad (6.5)$$

Next, we derive several corollaries of assumption (6.5).

Corollary 6.2. *Under the assumption (6.5), for all $a < b$,*

$$\mathbb{P} \left[\overline{\{0\} \times [8n, 12n)} \text{ is connected to } \overline{\{43n - 1\} \times [0, 4n)} \text{ in } T_{ab} \right] < \frac{c^9}{10^7}. \quad (6.6)$$

Proof of Corollary 6.2. Using reflectional symmetry and Corollary 4.3,

$$\begin{aligned} & \mathbb{P} \left[\overline{\{0\} \times [8n, 12n)} \text{ is connected to } \overline{\{43n - 1\} \times [0, 4n)} \text{ in } T_{ab} \right]^2 \\ &= \mathbb{P} \left[\overline{\{0\} \times [8n, 12n)} \text{ is connected to } \overline{\{43n - 1\} \times [0, 4n)} \text{ in } T_{ab} \right] \\ & \quad \cdot \mathbb{P} \left[\overline{\{0\} \times [0, 4n)} \text{ is connected to } \overline{\{43n - 1\} \times [8n, 12n)} \text{ in } T_{ab} \right] \\ &\leq c_*^{-1} \cdot \mathbb{P} \left[\overline{\{0\} \times [0, 4n)} \text{ is connected to } \overline{\{0\} \times [8n, 12n)} \text{ in } T_{ab} \right] \stackrel{(6.5)}{<} \frac{c^{18}}{10^{14}}. \end{aligned}$$




Figure 5: (a) illustration of the event in (6.6), (b) proof of Corollary 6.2.

□

Corollary 6.3. *Under the assumption (6.5), for all $a < b$,*

$$\mathbb{P} \left[\begin{array}{l} \text{there exist a simple path } \gamma \text{ from } \overline{\{0\} \times [0, 4n)} \text{ to } \overline{\{43n-1\} \times [0, 4n)} \text{ in } T_{ab} \\ \text{and a path } \gamma' \text{ from } \overline{\{0\} \times [8n, 12n)} \text{ in } T_{ab}, \text{ such that} \\ \text{the distance between } \bar{\gamma} \text{ and } \bar{\gamma}' \text{ is } \leq 2 \end{array} \right] < \frac{3 \cdot c^9}{10^7}. \quad (6.7)$$

In particular,

$$\mathbb{P} \left[\begin{array}{l} \text{there exist a simple path } \gamma \text{ from } \overline{\{0\} \times [0, 4n)} \text{ to } \overline{\{43n-1\} \times [0, 4n)} \text{ in } T \\ \text{and a path } \gamma' \text{ from } \overline{\{0\} \times [8n, 12n)} \text{ in } T, \text{ such that} \\ \text{the distance between } \bar{\gamma} \text{ and } \bar{\gamma}' \text{ is } \leq 2 \end{array} \right] \leq \frac{3 \cdot c^9}{10^7}. \quad (6.8)$$




Figure 6: An illustration of the event in (6.7).

Proof of Corollary 6.3. It suffices to prove (6.7), as (6.8) follows from (6.7) by sending $a \rightarrow -\infty$ and $b \rightarrow +\infty$.

Denote the event in (6.7) by A . By the total probability formula,

$$\begin{aligned} \mathbb{P}[A] \leq & \mathbb{P} \left[\overline{\{0\} \times [8n, 12n]} \text{ is connected to } \overline{\{0\} \times [0, 4n]} \text{ in } T_{ab} \right] \\ & + \mathbb{P} \left[\overline{\{0\} \times [8n, 12n]} \text{ is connected to } \overline{\{43n-1\} \times [0, 4n]} \text{ in } T_{ab} \right] \\ & + \mathbb{P} \left[\begin{array}{l} A, \overline{\{0\} \times [8n, 12n]} \text{ is not connected to } \overline{\{0\} \times [0, 4n]} \text{ in } T_{ab}, \\ \overline{\{0\} \times [8n, 12n]} \text{ is not connected to } \overline{\{43n-1\} \times [0, 4n]} \text{ in } T_{ab} \end{array} \right]. \end{aligned}$$

The sum of the first two probabilities is $< \frac{c_* \cdot c^{18}}{10^{14}} + \frac{c^9}{10^7}$, by the assumption (6.5) and (6.6).

Denote by A' the event in the third probability. For a configuration ω , let $P(\omega)$ be the set of vertices, which belong to at least one self-avoiding path from $\overline{\{0\} \times [0, 4n]}$ to $\overline{\{43n-1\} \times [0, 4n]}$ in T_{ab} , one may call it a backbone. Consider a local modification map f from A' to the event

$$A'' = \left\{ \omega'' : \begin{array}{l} \text{there exists a unique } z(\omega'') \in P(\omega'') \text{ connected to } \overline{\{0\} \times [8n, 12n]} \\ \text{by an open path contained in } T_{ab} \setminus P(\omega'') \text{ except for the vertex } z(\omega'') \end{array} \right\}$$

such that for all $\omega' \in A'$ and all $e \notin \overline{z(f(\omega')) + [-3, 3]^2 \times \{0\}^{d-2}}$, $f(\omega')_e = \omega'_e$. By Lemma 4.1, $\mathbb{P}[A'] \leq C_* \cdot \mathbb{P}[A''] \leq c_*^{-1} \cdot \mathbb{P}[A'']$. Since

$$A'' \subseteq \left\{ \overline{\{0\} \times [0, 4n]} \text{ is connected to } \overline{\{0\} \times [8n, 12n]} \text{ in } T_{ab} \right\},$$

we conclude that

$$\mathbb{P}[A'] \leq c_*^{-1} \cdot \mathbb{P} \left[\overline{\{0\} \times [0, 4n]} \text{ is connected to } \overline{\{0\} \times [8n, 12n]} \text{ in } T_{ab} \right] < \frac{c^{18}}{10^{14}},$$

where the last inequality follows from the assumption (6.5). Putting the bounds together,

$$\mathbb{P}[A] < \frac{c_* \cdot c^{18}}{10^{14}} + \frac{c^9}{10^7} + \frac{c^{18}}{10^{14}} \leq \frac{3 \cdot c^9}{10^7}.$$

□

Corollary 6.4. *Under the assumptions (6.4) and (6.5),*

$$\mathbb{P} \left[\begin{array}{l} \text{there exist a path } \gamma' \text{ from } \overline{\{0\} \times [0, 4n]} \text{ in } T \\ \text{and a path } \gamma'' \text{ from } \overline{\{0\} \times [16n, 20n]} \text{ in } T, \text{ such that} \\ \text{the distance between } \gamma' \text{ and } \gamma'' \text{ is } \leq 4 \end{array} \right] \leq \frac{12 \cdot c^8}{10^7}. \quad (6.9)$$




Figure 7: An illustration of the event in (6.9).

Proof of Corollary 6.4. Denote the event in (6.9) by A . By assumption (6.4),

$$\mathbb{P}\left[\overline{\{0\} \times [8n, 12n)} \text{ is connected to } \overline{\{43n-1\} \times [8n, 12n)} \text{ in } T\right] \geq c - 2 \frac{c}{10} \geq \frac{c}{2}.$$

Since the above event and the event A are increasing, by the FKG inequality,

$$\mathbb{P}[A] \leq \frac{2}{c} \cdot \mathbb{P}\left[A, \overline{\{0\} \times [8n, 12n)} \text{ is connected to } \overline{\{43n-1\} \times [8n, 12n)} \text{ in } T\right].$$

The intersection of the two events on the right hand side implies that for any path γ from $\overline{\{0\} \times [8n, 12n)}$ to $\overline{\{43n-1\} \times [8n, 12n)}$ in T , the distance from $\bar{\gamma}$ to $\bar{\gamma}' \cup \bar{\gamma}''$ is ≤ 2 . Thus, by (6.8),

$$\mathbb{P}[A] \leq \frac{2}{c} \cdot 2 \frac{3 \cdot c^9}{10^7} = \frac{12 \cdot c^8}{10^7}.$$

□

Step 4. The aim of this step is to introduce a certain event of positive probability, see Proposition 6.5. Our choice of this event will be clarified in Step 5.

Recall the definition of S from (6.3). For a configuration ω , let $C_S = C_S(\omega)$ be the set of all $z \in T$ connected to S by an open path in T . Let

$$f(\omega) = \mathbb{P}\left[\overline{\{0\} \times [4n, 8n)} \text{ is connected to } \overline{\{43n-1\} \times \mathbb{Z}} \text{ in } T \setminus \overline{C_S(\omega)} \mid C_S(\omega)\right],$$

and

$$g(\omega) = \mathbb{P}\left[\begin{array}{l} \text{there exists a path } \gamma' \text{ from } \overline{\{0\} \times [4n, 8n)} \text{ in } T, \text{ such that} \\ \text{the distance between } \bar{\gamma}' \text{ and } \overline{C_S(\omega)} \text{ is } \leq 4 \end{array} \mid C_S(\omega)\right].$$

We consider the following events:

$$\begin{aligned} A_1 &= \left\{ S \text{ is connected to } \overline{[0, 43n) \times \{2n\}} \text{ in } T \right\}, \\ A_2 &= \left\{ f(\omega) \geq \frac{c^2}{10} \right\}, \\ A_3 &= \left\{ g(\omega) \leq \frac{c^4}{1000} \right\}. \end{aligned}$$

Proposition 6.5. *Under the assumptions (6.2), (6.4), and (6.5),*

$$\mathbb{P}[A_1 \cap A_2 \cap A_3] \geq \frac{c^4}{10^3}.$$

Proof of Proposition 6.5. By assumptions (6.2) and (6.4),

$$\mathbb{P}[A_1] \geq c - \frac{c}{10} - \frac{c}{100} \geq \frac{c}{2}.$$

By the Markov inequality and (6.9),

$$\mathbb{P}[A_3^c] \leq \frac{1000}{c^4} \cdot \mathbb{E}[g] < \frac{1000}{c^4} \cdot \frac{12 \cdot c^8}{10^7} = \frac{12 \cdot c^4}{10^4}.$$

To bound $\mathbb{P}[A_1 \cap A_2]$ from below we use the Paley-Zygmund inequality. Using (6.9) and the FKG inequality, we first estimate

$$\begin{aligned}
\mathbb{E}[f(\omega) \cdot \mathbf{1}_{A_1}] &= \mathbb{P} \left[\begin{array}{l} S \text{ is connected to } \overline{[0, 43n) \times \{2n\}} \text{ in } T, \text{ and} \\ \overline{\{0\} \times [4n, 8n)} \text{ is connected to } \overline{\{43n-1\} \times \mathbb{Z}} \text{ in } T \setminus \overline{C_S(\omega)} \end{array} \right] \\
&\geq \mathbb{P} \left[\begin{array}{l} S \text{ is connected to } \overline{[0, 43n) \times \{2n\}} \text{ in } T, \text{ and} \\ \overline{\{0\} \times [4n, 8n)} \text{ is connected to } \overline{\{43n-1\} \times \mathbb{Z}} \text{ in } T \end{array} \right] - \frac{12 \cdot c^8}{10^7} \\
&\geq \mathbb{P}[A_1] \cdot \mathbb{P} \left[\overline{\{0\} \times [4n, 8n)} \text{ is connected to } \overline{\{43n-1\} \times \mathbb{Z}} \text{ in } T \right] - \frac{12 \cdot c^8}{10^7} \\
&\geq \frac{c}{2} \left(c - \frac{c}{10} \right) - \frac{12 \cdot c^8}{10^7} \geq \frac{c^2}{5}.
\end{aligned}$$

The Paley-Zygmund inequality for non-negative random variable X states that $\mathbf{P}[X \geq \frac{1}{2}\mathbf{E}[X]] \geq \frac{1}{4} \frac{(\mathbf{E}[X])^2}{\mathbf{E}[X^2]}$. We apply it to the measure $\mathbf{P}[\cdot] = \mathbb{E} \left[\mathbf{1}_{\cdot} \cdot \frac{\mathbf{1}_{A_1}}{\mathbb{P}[A_1]} \right]$, to get

$$\mathbb{E} \left[\mathbf{1}_{f(\omega) \geq \frac{1}{2} \cdot \mathbb{E}[f(\omega) \cdot \frac{\mathbf{1}_{A_1}}{\mathbb{P}[A_1]}]} \cdot \frac{\mathbf{1}_{A_1}}{\mathbb{P}[A_1]} \right] \geq \frac{1}{4} \cdot \left(\mathbb{E}[f(\omega) \cdot \frac{\mathbf{1}_{A_1}}{\mathbb{P}[A_1]}] \right)^2.$$

Thus,

$$\mathbb{P}[A_1 \cap A_2] \geq \frac{c^4}{100},$$

and we conclude that

$$\mathbb{P}[A_1 \cap A_2 \cap A_3] \geq \frac{c^4}{100} - \frac{12 \cdot c^4}{10^4} \geq \frac{c^4}{10^3}.$$

□

Step 5. We are ready to conclude. For a configuration ω , let $Q(\omega)$ be the set of vertices from T , which are connected to S by an open path in $\overline{[0, 43n) \times [2n, \infty)}$. Let $\Gamma(\omega)$ be the outer vertex boundary of $\overline{Q(\omega)}$, and $\Gamma'(\omega)$ the mirror reflection of Γ with respect to the hyperplane $\{x : x_2 = 2n - \frac{1}{2}\}$. We denote the connected component of $T \setminus (\Gamma \cup \Gamma')$ which contains 0 by V . Note that V is finite for any $\omega \in A_1$.

Let $X = \overline{\{0\} \times [4n, 8n)}$, and $X' = \overline{\{0\} \times [-4n, 0)}$. Note that X' is the mirror reflection of X with respect to the hyperplane $\{x : x_2 = 2n - \frac{1}{2}\}$. Moreover, if $\omega \in A_2 \cap A_3$, then both X and X' are contained in V .




Figure 8: An illustration of Γ , Γ' , and V for a configuration from the event $A_1 \cap A_2 \cap A_3$. Γ is the outer vertex boundary of the cluster of S in $[0, 43n) \times [2n, +\infty)$, Γ' is its mirror reflection with respect to the hyperplane $\{x : x_2 = 2n - \frac{1}{2}\}$, and V is the connected component of $T \setminus (\Gamma \cup \Gamma')$ containing the origin.

We consider an auxiliary probability space Ω' with configurations ω' and the same probability measure $\mathbb{P}$ on it. We compute

$$\begin{aligned}
& \mathbb{P} [X \text{ is connected to } X' \text{ in } T \text{ by an open path in } \omega'] \\
& \geq \mathbb{P} \otimes \mathbb{P} \left[(\omega, \omega') : \begin{array}{l} \omega \in A_1 \cap A_2 \cap A_3, \\ X \text{ is connected to } X' \text{ in } V(\omega) \text{ by an open path in } \omega' \end{array} \right] \\
& \stackrel{(*)}{\geq} C_*^{-1} \cdot \mathbb{P} \otimes \mathbb{P} \left[(\omega, \omega') : \begin{array}{l} \omega \in A_1 \cap A_2 \cap A_3, \\ X \text{ is not connected to } X' \text{ in } V(\omega) \text{ by an open path in } \omega', \\ X \text{ is connected to } \Gamma'(\omega) \text{ in } V(\omega) \text{ by an open path in } \omega', \\ X' \text{ is connected to } \Gamma(\omega) \text{ in } V(\omega) \text{ by an open path in } \omega', \\ \text{there is no open path } \pi \text{ in } \omega' \text{ from } X \text{ in } V(\omega) \\ \text{so that the distance between } \bar{\pi} \text{ and } \Gamma(\omega) \text{ is } \leq 4, \\ \text{there is no open path } \pi' \text{ in } \omega' \text{ from } X' \text{ in } V(\omega) \\ \text{so that the distance between } \bar{\pi}' \text{ and } \Gamma'(\omega) \text{ is } \leq 4 \end{array} \right] \\
& \geq C_*^{-1} \cdot \mathbb{E}_\omega \left[\mathbb{1}_{A_1 \cap A_2 \cap A_3}(\omega) \cdot \mathbb{P}_{\omega'} \left[\begin{array}{l} X \text{ is connected to } \Gamma'(\omega) \text{ in } V(\omega) \text{ by an open path in } \omega', \\ X' \text{ is connected to } \Gamma(\omega) \text{ in } V(\omega) \text{ by an open path in } \omega' \end{array} \right] \right] \\
& - C_*^{-1} \cdot \mathbb{E}_\omega \left[\mathbb{1}_{A_1 \cap A_2 \cap A_3}(\omega) \cdot \mathbb{P}_{\omega'} \left[\begin{array}{l} \text{there is an open path } \pi \text{ in } \omega' \text{ from } X \text{ in } V(\omega) \\ \text{so that the distance between } \bar{\pi} \text{ and } \Gamma(\omega) \text{ is } \leq 4, \\ \text{or} \\ \text{there is an open path } \pi' \text{ in } \omega' \text{ from } X' \text{ in } V(\omega) \\ \text{so that the distance between } \bar{\pi}' \text{ and } \Gamma'(\omega) \text{ is } \leq 4 \end{array} \right] \right] \\
& - C_*^{-1} \cdot \mathbb{P} [X \text{ is connected to } X' \text{ in } T \text{ by an open path in } \omega'] \\
& \geq C_*^{-1} \cdot \mathbb{E}_\omega [\mathbb{1}_{A_1 \cap A_2 \cap A_3}(\omega) \cdot [f(\omega)^2 - 2g(\omega)]] \\
& - C_*^{-1} \cdot \mathbb{P} [X \text{ is connected to } X' \text{ in } T \text{ by an open path in } \omega'].
\end{aligned}$$

The inequality (*) follows from Lemma 4.1 and a similar local transformation as in the proof of Corollary 6.3. (Mind that every path from X to Γ' in V and every path from X' to Γ have intersecting projections, and all the “intersection points” are sufficiently far away from $\Gamma \cup \Gamma'$ to allow for a local modification far away from $\Gamma \cup \Gamma'$.) The last inequality comes from the FKG inequality and the definitions of event A_1 and functions f and g .

By the definition of events A_2 and A_3 and Proposition 6.5,

$$\mathbb{P}[X \text{ is connected to } X' \text{ in } T] \geq c_* \cdot \left(\frac{c^4}{100} - 2 \cdot \frac{c^4}{1000} \right) \cdot \frac{c^4}{10^3}.$$

In particular, there exist $a < b$ such that

$$\mathbb{P}[X \text{ is connected to } X' \text{ in } T_{ab}] \geq \frac{c_* \cdot c^8}{10^6}.$$

From this we conclude, as in the argument of Step 3, that $p(44n, 43n) \geq c'$ (or simply observe that the above inequality contradicts the assumption (6.5)).

The proof of Proposition 6.1 is complete. □

References

- [1] M. Aizenman (1997) On the number of incipient spanning clusters. *Nucl. Phys. B* **485**, 551–582.
- [2] B. Bollobas and O. Riordan (2006) The critical probability for random Voronoi percolation in the plane is $\frac{1}{2}$. *Probab. Theory Related Fields* **136(3)**, 417–468.
- [3] H. Duminil-Copin, C. Hongler, and P. Nolin (2011) Connection probabilities and RSW-type bounds for the two-dimensional FK Ising model. *Comm. Pure Appl. Math.* **64(9)**, 1165–1198.
- [4] H. Duminil-Copin, V. Sidoravicius, and V. Tassion (2014) Absence of infinite cluster for critical Bernoulli percolation on slabs. *To appear in Comm. Pure Appl. Math.* arXiv:1401.7130.
- [5] H. Duminil-Copin, V. Sidoravicius, and V. Tassion (2015) Continuity of the phase transition for planar random-cluster and Potts models with $1 \leq q \leq 4$. arXiv:1505.04159.
- [6] G. R. Grimmett (1999) *Percolation*. Vol. 321. Springer Verlag.
- [7] H. Kesten (1982) *Percolation theory for mathematicians*. Birkhäuser Boston, Mass.
- [8] Ch. Newman, V. Tassion, and W. Wu (2015) Critical percolation and the minimal spanning tree in slabs. arXiv:1512.09107.
- [9] L. Russo (1978) A note on percolation. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **43(1)**, 39–48.
- [10] L. Russo (1981) On the critical percolation probabilities. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete* **56(2)**, 229–237.
- [11] P. D. Seymour and D. J. A. Welsh (1978) Percolation probabilities on the square lattice. *Annals of Discrete Mathematics* **3**, 227–245.
- [12] V. Tassion (2014) Crossing probabilities for Voronoi percolation. *To appear in Annals of Probability*, arXiv:1410.6773.